

**POTENTIAL HEALTH EFFECTS WITHIN OCCUPATIONAL
EXPOSURE TO ANESTHETICS OF HEALTH CARE WORKERS**

Eugenia Dănulescu¹, Mirela Ghițescu¹, Daniela Constantinescu¹, Carmen Cozmei¹,
Doina Havârneanu¹, Felicia Gradinariu¹, Irina Alexandrescu¹, V Cazuc¹,
Doina Ivanovici², R Dănulescu³, Raluca Dănulescu³

1. Institute of Public Health Iași, România
2. Authority of Public Health Iași, România
3. University of Medicine and Pharmacy Iași, România

Abstract. Aim. To assess respiratory system response in long time low-level occupational exposure to anesthetics (AE) and to identify appropriate prophylactic measures. **Materials and Methods.** Cross-section study on 128 health care workers occupationally exposed to AE (halothane, enflurane, isoflurane) vs matched controls. GC exposure evaluation, clinical examination, immunochemistry, respiratory function tests and oxidative stress markers (SOD, TBARS) were performed. To assess the possible effect of AE on the immune function, peripheral lymphocyte subpopulations, NK cells percentage and NK cytotoxic activity were determined in a subgroup (n=33). **Results.** AE concentrations in workplaces air were under but close to TLV. Irritative respiratory symptoms prevailed. Discrete distal obstructions were found in 45.3% of workers vs 22.8% in controls ($\chi^2=10.1$, $p<0.01$, OR=2.7, CI:1.4-5.2). SOD and TBARS were significantly higher in exposed ($p<0.01$). Serum immunoglobulins changes suggested alteration of the humoral immune response. The anaesthesiologist subgroup had a slightly reduced percentage of specific cytotoxicity (12.34%) versus exposed (16.06%). **Conclusions.** It seems that under TLV occupational exposure to anaesthetic gases could interfere with defence mechanisms. Further research could sustain the reconsideration of exposure standards - a suitable prophylactic approach.

Key words: respiratory health effects, occupational exposure, anesthetics, exposure standards

Rezumat. Scop. Evaluarea răspunsului aparatului respirator în expunerea ocupațională la aneestezice (AE) de nivele reduse, pe termen lung și identificarea măsurilor profilactice potrivite. **Material și metodă.** Studiu transversal pe 128 subiecți – personal medical expus ocupațional la AE (halotan, enfluran, izofluran) vs martori adecvați. S-au realizat: evaluarea expunerii prin gaz-cromatografie, examen clinic, probe funcționale ventilatorii, teste imunochimice și martori de stres oxidativ (superoxid dismutaza, radicali liberi). Pentru a evalua posibilul efect al AE asupra funcției imune, s-au determinat, în cadrul unui subgrup (n=33), subpopulațiile limfocitare periferice, procentajul și activitatea citotoxică a limfocitelor NK. **Rezultate.** Concentrațiile AE în aerul locurilor de muncă au atins nivele apropiate de limitele admisibile, fără însă a le depăși. Examenul clinic a arătat că simptomele iritative respiratorii au fost cele mai frecvente. S-au găsit obstrucții ventilatorii distale discrete la 45,3% dintre persoanele expuse din personalul medical față de 22,8% la lotul martor ($\chi^2=10,1$, $p<0.01$, OR=2,7, CI95%=1,4-5,2). Superoxid dismutaza și radicalii liberi au fost semnificativ mai mari la lotul expus ($p<0,05$). Modificările imunoglobulinelor serice au sugerat modificarea răspunsului umoral imun. Subgrupul de anesteziști a avut procentaje ușor

mai mici ale citotoxicității specifice (12,34%) față de lotul expus (16,06%). **Concluzii.** Se pare că expunerea ocupațională la gaze anestezice, chiar la nivele sub limitele admisibile, poate interfera cu mecanismele de apărare. Studii suplimentare viitoare ar putea motiva necesitatea reconsiderării standardelor de expunere. Este o abordare profilactică ce poate fi considerată ca adecvată în această situație.

Cuvinte cheie: efecte respiratorii, expunere ocupațională, anestezice, standarde de expunere

INTRODUCTION

Some data in scientific literature pointed out possibility that occupational exposure to anesthetics (AE) could interfere with health status of the involved health care workers (1, 3, 5, 12, 16).

The aim of the study was to assess respiratory system response in long-term low-level occupational exposure to volatile AE and to identify appropriate prophylactic measures (14).

The study hypothesis states that long-term low-level occupational exposure to frequently used volatile AE for general anesthesia during surgery could impair the health status of health care workers.

An epidemiological study on 128 health care workers (anesthesiology personnel and surgeons) occupationally exposed to volatile AE (halothane, enflurane, isoflurane) vs 87 matched controls was performed.

The exposed personnel and the controls were comparable concerning age (38.6 vs 37.7 years) and length of service (12.3 vs 11.9 years) homogeneity and distribution. Both lots were also comparable for other potential confounding factors (smoking, drug and alcohol consumption, dietary habits, extra-occupational exposure to irritative substances). In both lots people have jobs with similar physical solicitation; they live in the same town, in similar environments.

MATERIALS AND METHODS

Considering the relevant data in literature, exposure evaluation of volatile AE (halothane, enflurane, isoflurane) has been done by GC in the air of each specific workplace (operating rooms, intensive care units) in order to quantify as well as possible the exposure for every workplace (6, 8, 13, 15). Microclimate measurements were aimed to better assess the AE potential absorption degree.

The detailed investigation of health status comprised careful occupational and medical anamnesis, clinical examination, biochemical and immunochemical investigations (immunoglobulins A, G, E, and M), respiratory function tests. In order to test the genotoxic potential of AE occupational exposure we performed a cytogenetic investigation: micronuclei (MN) test in oral mucosa exfoliated cells (4, 9).

Oxidative stress markers have also been done: whole blood superoxide dismutase (SOD) and Thiobarbituric Acid Reactive Substances (TBARS) (7).

To assess the possible effect of AE on the immune function, peripheral lymphocyte subpopulations, NK cells percentage and NK cytotoxic activity were determined in a subgroup (n = 33 anesthesiologists) (11).

POTENTIAL HEALTH EFFECTS WITHIN OCCUPATIONAL EXPOSURE

RESULTS

Exposure evaluation of AE revealed that concentrations in workplaces air were under but close to TLV (fig. 1). The investigation of microclimate revealed conditions (especially high temperatures and low ventilation efficiency) which could facilitate the respiratory absorption. The health status investigation revealed irritative respiratory symptoms, allergic

conditions (dermatitis, rhinitis, laryngitis, asthma, conjunctivitis), cardiovascular changes, including arterial hypertension and EKG changes (arrhythmias, conduction disturbances, ischemia), arthropathies and musculoskeletal disorders, digestive apparatus disorders (chronic hepatitis, duodenal and gastric ulcers, cholecystitis). Prevalences are shown in figure 2.




Fig.1. Measured concentrations of volatile anesthetics (ppm)




Fig. 2. The prevalence of clinical entities (%)

From clinical point of view we found that irritative respiratory symptoms prevailed in exposed health care workers (33.6%), with a significant difference versus controls (Yates' chi

square = 8.41, $p=0.0037$, OR=2.88, 95% CI: 1.37-6.13). We also found rather frequent headaches, dizziness and nausea.




Fig. 3. The distribution of respiratory clinical entities

Considering the respiratory clinical entities (25.8% in exposed, vs 13.8% in controls - Yates' chi square = 3.8, $p=0.05$, OR=2.17, 95% CI: 1.00-4.80), we found acute rhinitis, chronic rhinitis, acute and chronic laryngitis, acute and chronic bronchitis and even asthma (fig. 3).

Spirometric investigations revealed that 52.34% of the exposed versus 28.74% of the controls had obstructive dysfunction (Yates' chi square = 10.85, $p = 0.0009$, OR = 2.72, 95% CI: 1.47-5.08).

Discrete distal obstructions were found in 45.3% of workers vs 22.8% in controls (Yates' chi square =10.1, $p=0.0013$, OR=2.7, 95%CI: 1.4-5.2). The distribution of obstructive dysfunction is shown in table 1.

We have found a medium correlation ($R = 0.44$, $p = 0.001$) between the obstructive syndromes and the presence of the respiratory irritative symptoms. Also a slight correlation ($R = 0.38$, $p=0.016$) was found between the obstructive syndromes and the intensity of occupational exposure (fig. 4).

POTENTIAL HEALTH EFFECTS WITHIN OCCUPATIONAL EXPOSURE

Table 1. Obstructive syndromes in exposed *versus* controls

	Normal subjects	Obstructions			
		Total obstructions	Distal Obstructive Syndrome FEV1 > 80% FEV1/VC > 80% FEF25-75%,MEF50%, MEF25%, PEF, MEF75% < 80%	Slight obstructions FEV1 < 80% FEV1/VC < 80%	Moderate and severe obstructions FEV1 < 55%
Exposed workers	61	67 52.34%	58 45.31%	7 5.47%	2 1.56%
Controls	62	25 28.74%	20 22.89%	5 5.74%	0 0%
Significance of difference		p=0.0013	p=0.0006	p=0.46	p=0.12




Fig. 4. Main findings in the occupational exposure to volatile anaesthetics.

Abnormal high immunoglobulins follow (fig. 5):
 values were found in exposed as




Fig. 5. Abnormal high immunoglobulins in exposed

The differences vs control were significant for IgE and IgM ($p < 0.05$) Ig A level correlated with Ig M and Ig G ($r = 0.34$, $p = 0.001$, $r = 0.31$, $p = 0.003$). Serum immunoglobulins changes suggested an alteration of the humoral immune response.

The blood oxidative balance was affected as shown by raised SOD activity in 55.2 % of exposed versus

10.1 % in controls and TBARS serum level raised above the upper reference limit in 83.1 % of exposed compared to 11.5 % in controls (fig. 6).

SOD and TBARS were also significantly higher in exposed (t test = 2.9, $p = 0.004$, respectively $t = 3.3$, $p = 0.001$) and slightly correlated with intensity of exposure ($R = 0.30$, $p = .051$).




Fig. 6. Oxidative stress markers in exposed vs control

POTENTIAL HEALTH EFFECTS WITHIN OCCUPATIONAL EXPOSURE

The cytogenetical investigation has shown that micronuclei (MN) in oral mucosa exfoliated cells were more

frequent in exposed (9.4% vs 2.3%, chi square = 4.26, $p=0.04$, OR=4.4 95%CI: 0.9 - 29.2)- see figure 7 (a, b).




Fig. 7. MN in oral mucosa

The assessment of the immune function by studying the NK peripheral lymphocytes (numerical and cytotoxic aspects) revealed the following results:

- there were no significant differences concerning NK cells number in AE exposed personnel versus control;
- the percentage of specific cytotoxicity was slightly reduced in AE exposed personnel (12.34%) versus exposed (16.06%).

CONCLUSIONS

- Some of the multifactorial diseases found in health care workers could have a pathogenic connection with the specific stress, oversteering workloads and long time shifts. So these health disorders could be considered as occupational conditioned diseases and the working conditions could influence their evolution.

- It seems that under TLV occupational exposure to anaesthetic gases could interfere with defence mechanisms.
- We found frequent irritative respiratory symptoms associated with obstructive dysfunction (mainly distal obstructive syndrome), with serum immunoglobulins changes that suggested an alteration of the humoral immune response, and also associated with an increase of oxidative stress markers. These data correlate with some findings of other researchers.
- The assessment of the possible effect of AE on the immune function suggests that anaesthetic gases could interfere with the immune function of the occupationally exposed subjects.
- Our findings seem to support the necessity of closely survey the immune status of exposed anaesthetists,

as well as the other biological modified parameters in order to completely understand their underlying action mechanisms.

- Further research could sustain the reconsideration of exposure standards, which seems to be a suitable prophylactic approach, especially considering some other European and international recommendations (2, 17).

REFERENCES

- Alexander BH, Checkoway H, Nagahama SI, Domino KB: *Cause-specific mortality risks of anesthesiologists*. *Anesthesiology*, 2000 Oct; 93 (4): 919-21.
- Badgwell Michael: *An evaluation of air safety source-control technology for the post anesthesia care unit*, *J PeriAnesthesia Nursing*, Volume 11, Issue 4, August 1996, Pages 207-222.
- Burm AGL: *Occupational hazards of inhalational anaesthetics*, *Best Practice & Research Clinical Anaesthesiology*, Volume 17, Issue 1, March 2003, Pages 147-161.
- Gunther W, Klaus H, Klaus S, Pawel S, Marion H, Hugo WR: *High-Level, but Not Low-Level, Occupational Exposure to Inhaled Anesthetics Is Associated with Genotoxicity in the Micronucleus Assay*, *Anesth. Analg.* 2001; 92:118-22.
- Hagmar L: *Is the life expectancy of anesthesiologists decreased?* *Scand J Work Environ Health* 2003; 29(2): 83-84.
- Hobbhahn J, Hoerauf K, Wiesner G, Schröegendorfer K, Taeger K: *Waste gas exposure during desflurane and isoflurane anaesthesia*. *Acta Anaesthesiol Scand* 1998; 42: 864-7.
- Hülya T, Ahmet A, Ahmet S: *Effect of Volatile Anesthetics on Oxidative Stress Due to Occupational Exposure*, *World Journal of Surgery*, Springer New York, ISSN 0364-2313, Issue Volume 29, Number 4 / April, 2005, Pages 540-542.
- Imberti R, Preseglio I, Imbriani M, Ghittori S, Cimino F, Mapelli A: *Low flow anaesthesia reduces occupational exposure to inhalation anaesthetics. Environmental and biological measurements in operating room personnel*. *Acta Anaesthesiol Scand* 1995; 39: 586-91.
- Jayakaran F, Thomas IM: *Sister-chromatid exchanges in anesthetists*. *Indian J Occup Environ Med* 2005; 9: 86-89.
- Karakaya A, Tuncel N, Yücesoy B, et al: *The effects of volatile anaesthetic agents on human immune system function via occupational exposure*. *Immunopharmacol Immunotoxicol* 1992; 14: 251-9.
- Karelová J, Jablonická A, Gavora J, Hano L: *Chromosome and sister-chromatid exchange analysis in peripheral lymphocytes, and mutagenicity of urine in anesthesiology personnel*. *Int Arch Occup Environ Health* 1992; 64: 303-6.
- Katz JD: *Do anesthesiologists die at a younger age than other physicians? Age-adjusted death rates*. *Anesth Analg.* 2004 Apr; 98(4): 1111-3.
- Marx T, Schmidt M, Schirmer U, Reinelt H: *Pollution of the Environment and the Workplace with Anesthetic Gases*; *International Anesthesiology Clinics*. 2001, 39(2): 15-27.
- Mierdl S, Byhahn C, Abdel-Rahman U, Matheis G, Westphal K: *Occupational exposure to inhalational anesthetics during cardiac surgery on cardiopulmonary bypass*, *The Annals of Thoracic Surgery*, Volume 75, Issue 6, June 2003, Pages 1924-1927.
- Morgan GEdward Jr, Mikhail SM, Murray JM: *Clinical Anesthesiology*,

POTENTIAL HEALTH EFFECTS WITHIN OCCUPATIONAL EXPOSURE

- 4th Edition, 2005, McGraw-Hill Medical; ISBN 0-07-142358-3, Section V. Special Problems, Occupational Hazards in Anesthesiology.
16. Murat O, Sibel B, Deniz K, Serhat K, Ayla T: *Behavioural effects of chronic exposure to subanesthetic concentrations of halothane, sevoflurane and desflurane in rats*, Canadian Journal of Anesthesia 2006, 53: 653-658.
 17. Unceta-Barrenechea OB, Vicinay Pinedo S, Garran Sabando B, Serna de Andres A, Seoane de Lucas A: *Occupational exposure of the anesthesiologist to nitrous oxide and halothane. Control measures*, Rev Esp Anesthesiol Reanim. 1989 Sep-Oct; 36(5): 267-75.